arXiv:1212.5839v1 [math.DG] 23 Dec 2012

Slant curves in 3-dimensional normal almost paracontact metric manifolds

Joanna Wełyczko

December 27, 2012

Abstract

The presented paper is devoted to study the curvature and torsion of slant Frenet curves in 3-dimensional normal almost paracontact metric manifolds. Moreover, in this class of manifolds, properties of non-Frenet slant curves (with null tangents or null normals) are obtained. The achieved results are illustrated by examples.

1 Preliminaries

Let M be a $(2n+1)$-dimensional differentiable manifold. Let φ be a $(1,1)$-tensor filed, ξ a vector filed and η a 1-form on M. Then (φ,ξ,η) is called an almost paracontac structure on M if

(i) $\eta(\xi)=1, \quad \varphi^2=\mathrm{Id}-\eta\otimes\xi$,

(ii) the tensor filed φ induces an almost paracomplex structure on the distribution $\mathcal{D}=ker\,\eta$, that is, the eigendistributions $\mathcal{D}^+$, $\mathcal{D}^-$ corresponding to the eigenvalues 1, -1 of φ, respectively, have equal dimension n.

M is said to be almost paracontact manifold if it is endowed with an almost paracontact structure (cf. [2, 10, 14, 16]).

Let M be an almost paracontact manifold. M will be called an almost paracontact metric manifold if it is additionally endowed with a pseudo-Riemannian metric g of signature $(n+1,n)$ and such that

$$g(\varphi X,\varphi Y)=\,-\,g(X,Y)+\eta(X)\eta(Y).$$

For such a manifold, we additionally have $\eta(X)=g(X,\xi)$, $\eta(\xi)=1$, $\varphi\xi=0$, $\eta\circ\varphi=0$. Moreover, we can define a skew-symmetric 2-form $\varPhi$ by $\varPhi(X,Y)=g(X,\varphi Y)$, which is called the fundamental form corresponding to the structure. Note that $\eta\wedge\Phi$ is up to a constant factor the Riemannian volume element of M.

On an almost paracontact manifold, one defines the $(2,1)$-tensor filed $N^{(1)}$ by

$$N^{(1)}(X,Y)=[\varphi,\,\varphi](X,Y)-2\,d\eta(X,Y)\xi,$$

where $[\varphi, \varphi]$ is the Nijenhuis torsion of φ given by

$$[\varphi,\, \varphi](X,Y) = \varphi^2[X,Y] + [\varphi X, \varphi Y] - \varphi[\varphi X, Y] - \varphi[X, \varphi Y].$$

If $N^{(1)}$ vanishes identically, then the almost paracontact manifold (structure) is said to be normal (cf. [14] and recent papers [2, 16]). The normality condition says that the almost paracomplex structure J defined on $M \times \mathbb{R}$ by

$$J\Big(X, \lambda\frac{d}{dt}\Big) = \Big(\varphi X + \lambda\xi, \eta(X)\frac{d}{dt}\Big)$$

is integrable (paracomplex).

Since that time, we will consider manifolds of dimension 3. The following theorem presents conditions equivalent to the normality of 3 dimensional almost paracontact manifold, which we will use later in the work.

Theorem 1.1 [15] *For a 3-dimensional almost paracontact metric manifold M, the following three conditions are mutually equivalent*

(a) M is normal,

(b) there exist functions α, β on M such that

$$(\nabla_X\varphi)Y = \beta(g(X,Y)\xi - \eta(Y)X) + \alpha(g(\varphi X, Y)\xi - \eta(Y)\varphi X), \tag{1.1}$$

(c) there exist functions α, β on M such that

$$\nabla_X\xi = \alpha(X - \eta(X)\xi) + \beta\varphi X. \tag{1.2}$$

Corollary 1.2 [15] *The functions α, β realizing (1.1) as well as (1.2) are given by*

$$2\alpha = \mathrm{Trace}\,\{X \to \nabla_X\xi\}, \qquad 2\beta = \mathrm{Trace}\,\{X \to\ \varphi\nabla_X\xi\}. \tag{1.3}$$

A 3-dimensional normal almost paracontact metric manifold is said to be

- paracosymplectic manifold if $\alpha = \beta = 0$, [6]
- quasi-para-Sasakian if and only if $\alpha = 0$ and $\beta \neq 0$, [10, 15]
- β-para-Sasakian if and only if $\alpha = 0$ and $\beta \neq 0$ and β is constant, in particular, para-Sasakian if $\beta = -1$ ([15, 16])
- α-para-Kenmotsu if $\alpha \neq 0$ and α is constant and $\beta = 0$.

At the end of this section, let $\gamma\colon I \to M$, I being an interval, be a curve in M such that $g(\dot\gamma, \dot\gamma) = \pm 1$ or $g(\dot\gamma, \dot\gamma) = 0$ and let $\nabla_{\dot\gamma}$ denote the covariant differentiation along γ.

We say that the curve γ is a slant curve in 3-dimensional normal almost paracontact metric manifold, if

$$g(\dot\gamma, \xi) = \eta(\dot\gamma) = c \text{ and } c \text{ is constant.}$$

In the particular case, if $c = 0$ the curve γ is called Legendre curve [15].

Remark 1.3 *In normal almost contact metric manifold with Riemannian metric g, the value of $g(\dot\gamma,\xi)$ satisfies $-1\leqslant g(\dot\gamma,\xi)\leqslant 1$, so that we can define the structural angel of γ i.e the function $\theta: I\to[0,2\pi)$ given by*

$$\cos\theta(t)=g(\dot\gamma(t),\xi)=\eta(\dot\gamma(t)).$$

Then, the curve γ is said to be a slant curve, (or θ-slant curve), if θ is a constant function, (see [3, 4, 5]*).*

Let $\gamma: I\to M$ be a slant curve in M such that $g(\dot\gamma,\dot\gamma)=\varepsilon_1=\pm1$. Let us consider tree vector fields $\dot\gamma,\varphi\dot\gamma,\xi$ for which, we have

$$g(\dot\gamma,\dot\gamma)=\varepsilon_1,\ g(\dot\gamma,\xi)=c,\ g(\varphi\dot\gamma,\varphi\dot\gamma)=-\varepsilon_1+c^2,\ g(\xi,\xi)=1,\ g(\dot\gamma,\varphi\dot\gamma)=g(\xi,\varphi\dot\gamma)=0.$$

This vector field are linearly independent and form the base of TM if and only if $\varepsilon_1-c^2\neq 0$. In this case, we define an orthonormal frame along γ as follows:

$$F_1=\dot\gamma,\quad F_2=\frac{\varphi\dot\gamma}{\sqrt{|\varepsilon_1-c^2|}},\quad F_3=\frac{\xi-\varepsilon_1 c\dot\gamma}{\sqrt{|\varepsilon_1-c^2|}},\tag{1.4}$$

where $g(F_1,F_1)=\varepsilon_1$, $g(F_2,F_2)=sgn\,(-(\varepsilon_1-c^2))=\upsilon$, $g(F_3,F_3)=-\varepsilon_1\upsilon$.

The frame (F_1,F_2,F_3) will play an auxiliary role in the proofs of Theorems 2.2 and 4.1. We additionally have

Proposition 1.4 *Let M be a 3-dimensional normal almost paracontact metric manifold. If $\gamma\colon I\to M$ is a slant Frenet curve of osculating order 3 which is not an integral curve of ξ, then*

$$\begin{aligned}
\nabla_{\dot\gamma}F_1 &= \upsilon\delta\sqrt{|\varepsilon_1-c^2|}F_2-\varepsilon_1\alpha\sqrt{|\varepsilon_1-c^2|}F_3, &(1.5)\\
\nabla_{\dot\gamma}F_2 &= -\varepsilon_1\delta\sqrt{|\varepsilon_1-c^2|}F_1+(\varepsilon_1\beta-\upsilon c\delta)F_3, &(1.6)\\
\nabla_{\dot\gamma}F_3 &= -\varepsilon_1\upsilon\alpha\sqrt{|\varepsilon_1-c^2|}F_1+(\beta-\varepsilon_1\upsilon c\delta)F_2, &(1.7)
\end{aligned}$$

where δ is a function defined by $\delta=\dfrac{g(\nabla_{\dot\gamma}\dot\gamma,\varphi\dot\gamma)}{|\varepsilon_1-c^2|}$, and for simplicity we write α, β instead of the composed functions $\alpha\circ\gamma$, $\beta\circ\gamma$ with α, β being the same as in (1.3).

Separately, we discuss the case when $\dot\gamma,\varphi\dot\gamma,\xi$ are linearly dependent. Let us note that $\dot\gamma,\varphi\dot\gamma,\xi$ are linearly dependent if and only if

$$\dot\gamma=c\xi\quad\text{or}\quad\dot\gamma=c\xi\pm\varphi\dot\gamma.\tag{1.8}$$

Indeed, if $\dot\gamma,\varphi\dot\gamma,\xi$ are linearly dependent then $\varepsilon_1=1$ and $c^2=1$. Hence $g(\varphi\dot\gamma,\varphi\dot\gamma)=0$ and either $\varphi\dot\gamma=0$ or $\varphi\dot\gamma$ is an isotropic vector field.

If $\varphi\dot\gamma=0$ then $0=\varphi^2\dot\gamma=\dot\gamma-c\xi$ and $\dot\gamma=c\xi$.

If $\varphi\dot\gamma$ is an isotropic vector field then, we can write $\dot\gamma=a\xi+b\varphi\dot\gamma$, for some functions a and b. Using $g(\dot\gamma,\dot\gamma)=1$ and $g(\dot\gamma,\xi)=c$, we find $\dot\gamma=c\xi+b\varphi\dot\gamma$. Now

$$\varphi\dot\gamma=b\varphi^2\dot\gamma=b(\dot\gamma-c\xi)=b(c\xi+b\varphi\dot\gamma-c\xi)=b^2\varphi\dot\gamma.$$

Hence $b^2=1$ and $\dot\gamma=c\xi\pm\varphi\dot\gamma$.

2 Frenet slant curves

Let M be a 3-dimensional pseudo-Riemannian manifold with metric g.

Let $\gamma\colon I \to M$, I being an interval, be a curve in M.

We say that γ is a Frenet curve if $g(\dot\gamma,\dot\gamma) = \varepsilon_1$, $\varepsilon_1 = \pm 1$ and one of the following three cases hold

(a) γ is of osculating order 1, i.e., $\nabla_{\dot\gamma}\dot\gamma = 0$ (geodesics);

(b) γ is of osculating order 2, i.e., there exist two orthonormal vector fields $E_1(=\dot\gamma)$, E_2, $(g(E_2,E_2) = \varepsilon_2 = \pm 1)$ and a positive function κ (the curvature) along γ such that

$$\nabla_{\dot\gamma}E_1 = \kappa\varepsilon_2 E_2, \quad \nabla_{\dot\gamma}E_2 = -\kappa\varepsilon_1 E_1;$$

(c) γ is of osculating order 3, i.e., there exist three orthonormal vector fields $E_1(=\dot\gamma)$, E_2, E_3, $(g(E_2,E_2) = \varepsilon_2 = \pm 1$, $g(E_3,E_3) = \varepsilon_3 = \pm 1)$ and two positive functions κ (the curvature) and τ (the torsion) along γ such that

$$\nabla_{\dot\gamma}E_1 = \kappa\varepsilon_2 E_2, \quad \nabla_{\dot\gamma}E_2 = -\kappa\varepsilon_1 E_1 + \tau\varepsilon_3 E_3, \quad \nabla_{\dot\gamma}E_3 = -\tau\varepsilon_2 E_2.$$

Proposition 2.1 *Let M be a 3-dimensional normal almost paracontact metric manifold. If $\gamma\colon I \to M$ is a slant Frenet curve in M, such that $g(\dot\gamma,\dot\gamma) = 1$ and $g(\dot\gamma,\xi) = c = \pm 1$ then γ is a geodesics.*

Proof In view of (1.8), we consider two opportunities.

If $\dot\gamma = c\xi$ then from (1.2) $\nabla_{\dot\gamma}\dot\gamma = \nabla_\xi\xi = 0$ and γ is a geodesics.

If $\dot\gamma = c\xi \pm \varphi\dot\gamma$ using (1.2) and (1.1), we get

$$\begin{aligned} \nabla_{\dot\gamma}\dot\gamma &= c\nabla_{\dot\gamma}\xi \pm \nabla_{\dot\gamma}\varphi\dot\gamma = c\nabla_{\dot\gamma}\xi \pm (\nabla_{\dot\gamma}\varphi)\dot\gamma \pm \varphi\nabla_{\dot\gamma}\dot\gamma \\ &= (\alpha \mp \beta)(c\dot\gamma - \xi \mp c\varphi\dot\gamma) \pm \varphi\nabla_{\dot\gamma}\dot\gamma = \pm\varphi\nabla_{\dot\gamma}\dot\gamma. \end{aligned}$$

Then $g(\nabla_{\dot\gamma}\dot\gamma, \nabla_{\dot\gamma}\dot\gamma) = \pm g(\nabla_{\dot\gamma}\dot\gamma, \varphi\nabla_{\dot\gamma}\dot\gamma) = 0$ and γ is a Frenet curve so that γ is a geodesics. □

Theorem 2.2 generalizes results obtained for the Legendre curve in [15].

Theorem 2.2 *Let M be a 3-dimensional normal almost paracontact metric manifold. If $\gamma\colon I \to M$ is a slant Frenet curve of osculating order 3 in M, then its curvature and torsion are given by*

$$\kappa = \sqrt{|\varepsilon_1 - c^2||\alpha^2 - \varepsilon_1\delta^2|}, \tag{2.1}$$

$$\tau = \Big| sgn(1 - \varepsilon_1 c^2)\beta + c\delta + \frac{\alpha\dot\delta - \dot\alpha\delta}{\alpha^2 - \varepsilon_1\delta^2}\Big|. \tag{2.2}$$

where δ is a function defined by $\delta = \dfrac{g(\nabla_{\dot\gamma}\dot\gamma, \varphi\dot\gamma)}{|\varepsilon_1 - c^2|}$, and for simplicity we write α, β instead of the composed functions $\alpha\circ\gamma$, $\beta\circ\gamma$ with α, β being the same as in (1.3).

Proof Let γ be a slant Frenet curve of osculating order 3 in M. From Proposition 2.1, we know, that for such a curve $\varepsilon_1 - c^2 \neq 0$ and so that we can use the frame (F_1, F_2, F_3) from (1.4) to calculate the curvature function and the torsion function of a curve γ.

We obtain the curvature function using (1.5) and calculating the length of $\nabla_{\dot\gamma}\dot\gamma$. We get

$$\kappa^2\varepsilon_2 = -\varepsilon_1\upsilon|\varepsilon_1 - c^2|(\alpha^2 - \varepsilon_1\delta^2)$$

so that the curvature is given by (2.1) and $\varepsilon_2 = sgn(-\varepsilon_1\upsilon(\alpha^2 - \varepsilon_1\delta^2)) = \pm 1$.

Using (1.5), (1.6) and (1.7), we find

$$E_2 = \frac{1}{\varepsilon_2\kappa}\nabla_{\dot\gamma}E_1 = \frac{\varepsilon_2(\upsilon\delta F_2 - \varepsilon_1\alpha F_3)}{\sqrt{|\alpha^2 - \varepsilon_1\delta^2|}},$$

Let us denote $p = \sqrt{|\alpha^2 - \varepsilon_1\delta^2|}$ and calculate

$$\begin{aligned}
\nabla_{\dot\gamma}E_2 &= \varepsilon_2\upsilon\dot\gamma\Big(\frac{\delta}{p}\Big)F_2 + \varepsilon_2\upsilon\Big(\frac{\delta}{p}\Big)\nabla_{\dot\gamma}F_2 \\
&\quad - \varepsilon_1\varepsilon_2\dot\gamma\Big(\frac{\alpha}{p}\Big)F_3 - \varepsilon_1\varepsilon_2\Big(\frac{\alpha}{p}\Big)\nabla_{\dot\gamma}F_3 \\
&= -\varepsilon_1\kappa E_1 + \Big(\varepsilon_2\upsilon\dot\gamma\Big(\frac{\delta}{p}\Big) - \varepsilon_2\frac{\alpha\upsilon}{p}\Big(\varepsilon_1\upsilon\beta - c\delta\Big)\Big)F_2 \\
&\quad + \Big(-\varepsilon_1\varepsilon_2\dot\gamma\Big(\frac{\alpha}{p}\Big) + \varepsilon_2\frac{\delta}{p}\Big(\varepsilon_1\upsilon\beta - c\delta\Big)\Big)F_3 \\
&= -\varepsilon_1\kappa E_1 + \varepsilon_2\Big(-\varepsilon_1\upsilon\beta + c\delta + \frac{\alpha\dot\delta - \dot\alpha\delta}{\alpha^2 - \varepsilon_1\delta^2}\Big)\frac{\upsilon\alpha F_2 - \delta F_3}{\sqrt{|\alpha^2 - \varepsilon_1\delta^2|}}.
\end{aligned} \tag{2.3}$$

Hence from (2.3), in view of Frenet equations, we have

$$\begin{aligned}
\varepsilon_3\tau^2 &= \upsilon\, sgn(\alpha^2 - \varepsilon_1\delta^2)\Big(-\varepsilon_1\upsilon\beta + c\delta + \frac{\alpha\dot\delta - \dot\alpha\delta}{\alpha^2 - \varepsilon_1\delta^2}\Big)^2 \\
&= -\varepsilon_1\varepsilon_2\Big(sgn(1 - \varepsilon_1c^2)\beta + c\delta + \frac{\alpha\dot\delta - \dot\alpha\delta}{\alpha^2 - \varepsilon_1\delta^2}\Big)^2.
\end{aligned}$$

This gives (2.2) and $\varepsilon_3 = -\varepsilon_1\varepsilon_2$. □

Below, we present the functions of curvature and torsion of a slant Frenet curves of osculating order 3 in some subclasses of 3-dimensional normal almost paracontact metric manifold.

Corollary 2.3 *Let M be a 3-dimensional manifold, $\gamma\colon I \to M$ be a slant Frenet curve of osculating order 3 in M.*

If M is a paracosymplectic manifold then the curvature and torsion of γ are given by

$$\kappa = \sqrt{|\varepsilon_1 - c^2|}|\delta|, \quad \tau = |c\delta|.$$

If M is a quasi-para-Sasakian manifold then the curvature and torsion of γ are given by

$$\kappa = \sqrt{|\varepsilon_1 - c^2||\delta|}, \quad \tau = \Big| sgn(1-\varepsilon_1 c^2)\beta + c\delta \Big|.$$

If M is a para-Sasakian manifold then the curvature and torsion of γ are given by

$$\kappa = \sqrt{|\varepsilon_1 - c^2||\delta|}, \quad \tau = \Big| sgn(1-\varepsilon_1 c^2) + c\delta \Big|.$$

If M is an α-Kenmotsu manifold then the curvature and torsion of γ are given by

$$\kappa = \sqrt{|\varepsilon_1 - c^2||\alpha^2 - \varepsilon_1\delta^2|}, \quad \tau = \left| c\delta + \frac{\alpha\dot{\delta}}{\alpha^2 - \varepsilon_1\delta^2} \right|.$$

Moreover δ is a function defined by $\delta = \dfrac{g(\nabla_{\dot\gamma}\dot\gamma, \varphi\dot\gamma)}{|\varepsilon_1 - c^2|}$, and for simplicity we write α, β instead of the composed functions $\alpha \circ \gamma$, $\beta \circ \gamma$ with α, β being the same as in (1.3).

3 Null slant Curves

Next, we consider the case when γ is a null curve, i.e has a null tangent ($g(\dot\gamma, \dot\gamma) = 0$).

We consider null curve γ which is not a geodesic, then $g(\nabla_{\dot\gamma}\dot\gamma, \nabla_{\dot\gamma}\dot\gamma) \neq 0$. We take a parametrization of γ such that $g(\nabla_{\dot\gamma}\dot\gamma, \nabla_{\dot\gamma}\dot\gamma) = 1$ and define (see [7, 8, 9, 13])

$$T = \dot\gamma, \quad N = \nabla_{\dot\gamma}T, \quad \tau = \frac{1}{2}g(\nabla_{\dot\gamma}N, \nabla_{\dot\gamma}N), \quad W = -\nabla_{\dot\gamma}N - \tau T, \tag{3.1}$$

where

$$g(T,W) = g(N,N) = 1 \quad \text{and} \quad g(T,T) = g(T,N) = g(W,W) = g(W,N) = 0.$$

Then, we obtain the Cartan equations for null curve γ.

$$\nabla_{\dot\gamma}T = N, \quad \nabla_{\dot\gamma}W = \tau N. \quad \nabla_{\dot\gamma}N = -\tau T - W.$$

In [15], we proved that every null Legendre curve in 3-dimensional normal almost paracontact metric manifold is a geodesic.

In this section, we consider a slant non-geodesic null curve $\gamma : I \to M$.

Theorem 3.1 *If $\gamma : I \to M$ is a slant non-geodesic null curve in a 3-dimensional normal almost paracontact metric manifold then*

$$\tau = -\frac{\alpha^2 c^2}{2} - \dot\alpha c \pm \beta - \frac{1}{2c^2}, \tag{3.2}$$

$$N = \alpha c\dot\gamma \pm \frac{1}{c}\varphi\dot\gamma, \tag{3.3}$$

$$W = \frac{-\alpha^2 c^2 - 1}{2c}\dot\gamma \mp \alpha\varphi\dot\gamma + \frac{1}{c^2}\xi, \tag{3.4}$$

where for simplicity we write α, β instead of the composed functions $\alpha\circ\gamma$, $\beta\circ\gamma$ with α, β being the same as in (1.3).

Proof First, we find a vector field N. Let $\varphi\dot\gamma = p\dot\gamma + qN + rW$ for some functions p, q, r. We find

$$r = g(\varphi\dot\gamma,\dot\gamma) = 0,\quad g(\varphi\dot\gamma,\varphi\dot\gamma) = c^2 = q^2,\quad g(\varphi\dot\gamma,\xi) = pc+q\alpha c^2 = c(p\pm\alpha c^2) = 0.$$

Hence $r = 0$, $q = \pm c$, $p = \mp\alpha c^2$ and $\varphi\dot\gamma = \mp\alpha c^2\dot\gamma \pm cN$, which leads to (3.3).

Now, applying (1.1) and (3.3), we calculate

$$\begin{aligned}\nabla_{\dot\gamma}N &= \dot\alpha c\dot\gamma + \alpha c\nabla_{\dot\gamma}\dot\gamma \pm \frac{1}{c}((\nabla_{\dot\gamma}\varphi)\dot\gamma + \varphi\nabla_{\dot\gamma}\dot\gamma) \\ &= \Big(\dot\alpha ca + \alpha^2c^2 \mp \beta + \frac{1}{c}\Big)\dot\gamma \pm \alpha\varphi\dot\gamma - \frac{1}{c}\xi.\end{aligned} \tag{3.5}$$

Next, in view of definition τ and W in (3.1), we use (3.5) and get (3.2) and (3.4). □

In the following corollary, we present the properties of slant null curves in some subclasses of 3-dimensional normal almost paracontact metric manifold.

Corollary 3.2 *Let M be a 3-dimensional manifold, $\gamma\colon I\to M$ be a slant non-geodesic null curve in M.*

If M is a paracosymplectic manifold then for γ, we have

$$\tau = -\frac{1}{2c^2},\quad N = \pm\frac{1}{c}\varphi\dot\gamma,\quad W = \frac{1}{c^2}\xi.$$

If M is a quasi-para-Sasakian manifold then for γ, we have

$$\tau = \pm\beta - \frac{1}{2c^2},\quad N = \pm\frac{1}{c}\varphi\dot\gamma,\quad W = \frac{1}{c^2}\xi.$$

If M is a para-Sasakian manifold then for γ, we have

$$\tau = \pm 1 - \frac{1}{2c^2},\quad N = \pm\frac{1}{c}\varphi\dot\gamma,\quad W = \frac{1}{c^2}\xi.$$

If M is an α-Kenmotsu manifold then for γ, we have

$$\tau = -\frac{\alpha^2c^2}{2} - \frac{1}{2c^2},\quad N = \alpha c\dot\gamma \pm \frac{1}{c}\varphi\dot\gamma,\quad W = \frac{-\alpha^2c^2-1}{2c}\dot\gamma \mp \alpha\varphi\dot\gamma + \frac{1}{c^2}\xi.$$

.

4 Slant curves with null normal

At last, we investigate curves with null normals (for such curves in Minkowski spaces $\mathbb{E}^3_1$, see among others [1, 11, 12]. We say $\gamma\colon I\to M$ is a curve with null normal if

$$g(\dot\gamma,\dot\gamma) = \varepsilon_1 = \pm1,\quad \nabla_{\dot\gamma}\dot\gamma \neq 0,\quad g(\nabla_{\dot\gamma}\dot\gamma,\nabla_{\dot\gamma}\dot\gamma) = 0.$$

For such a curve, we define

$$T = \dot{\gamma}, \quad N = \nabla_{\dot{\gamma}} T, \quad W,$$

where

$$\begin{array}{l} g(T,T) = \varepsilon_1, \quad g(N,W) = 1, \\ g(N,N) = g(W,W) = g(T,N) = g(T,W) = 0. \end{array} \tag{4.1}$$

The signature of metric g is $(-,+,+)$ so that $\varepsilon_1 = 1$.

We obtain the following Cartan equations for null normal curve γ. (In Minkowski 3-space see [12])

$$\nabla_{\dot{\gamma}} T = N, \quad \nabla_{\dot{\gamma}} N = \kappa N, \quad \nabla_{\dot{\gamma}} W = -T - \kappa W.$$

Theorem 4.1 *If $\gamma : I \to M$ is a slant non-geodesic curve with null normal in a 3-dimensional normal almost paracontact metric manifold, for which $c^2 \neq 1$ then $\alpha = \pm \dfrac{g(N, \varphi\dot{\gamma})}{1-c^2} \neq 0$, and*

$$N = -\alpha(\xi - c\dot{\gamma} \pm \varphi\dot{\gamma}), \tag{4.2}$$

$$\kappa = \frac{\dot{\alpha}}{\alpha} \pm \beta + \alpha c, \tag{4.3}$$

$$W = \frac{-1}{2\alpha(1-c^2)}(\xi - c\dot{\gamma} \mp \varphi\dot{\gamma}), \tag{4.4}$$

where for simplicity we write α, β instead of the composed functions $\alpha \circ \gamma$, $\beta \circ \gamma$ with α, β being the same as in (1.3).

Proof For a slant curve with null normal for which $c^2 \neq 1$, we can use an orthonormal frame along γ as in (1.4). We put

$$N = \nabla_{\dot{\gamma}}\dot{\gamma} = aF_2 + bF_3, \tag{4.5}$$

for some function a and b and calculate $g(N,N) = a^2 - b^2 = 0$. Hence $a = \pm b$. Moreover, using (1.2), we get

$$\begin{aligned} b = g(N, F_3) &= -\frac{1}{\sqrt{|1-c^2|}} g(\nabla_{\dot{\gamma}}\xi, \dot{\gamma}) \\ &= -\alpha\sqrt{|1-c^2|}. \end{aligned} \tag{4.6}$$

Hence $\alpha \neq 0$ and putting (4.6) to (4.5), we obtain (4.2).

To find the curvature, we use (1.2), (4.2) , (1.1) and calculate

$$\begin{aligned} \nabla_{\dot{\gamma}} N &= -\dot{\alpha}(\xi - c\dot{\gamma} \pm \varphi\dot{\gamma}) \\ &\quad -\alpha(\nabla_{\dot{\gamma}}\xi - c\nabla_{\dot{\gamma}}\dot{\gamma} \pm (\nabla_{\dot{\gamma}}\varphi)\dot{\gamma} \pm \varphi\nabla_{\dot{\gamma}}\dot{\gamma}) \\ &= (-\dot{\alpha} \mp \alpha\beta - \alpha^2 c)(\xi - c\dot{\gamma} \pm \varphi\dot{\gamma}) \\ &= \Big(\frac{\dot{\alpha}}{\alpha} \pm \beta + \alpha c\Big) N. \end{aligned}$$

In view of Cartan frame of γ, we obtain (4.3).

At least, we put $W = dF_1 + eF_2 + fF_3$ for some functions d, e, f and using (4.1), we get (4.4). $\square$

As an immediate consequence of the above theorem, we obtain

Theorem 4.2 *Let $\gamma : I \to M$ be a Legendre curve with null normal in a 3-dimensional normal almost paracontact metric manifold M. Then $\alpha = \pm g(N, \varphi\dot\gamma)$, $\alpha \neq 0$ and we respectively have*

$$T = \dot\gamma, \quad N = -\alpha(\xi \pm \varphi\dot\gamma), \quad W = \frac{-1}{2\alpha}(\xi \mp \varphi\dot\gamma) \quad \textit{and} \quad \kappa = \left(\frac{\dot\alpha}{\alpha} \pm \beta\right),$$

where for simplicity we write α, β instead of the composed functions $\alpha \circ \gamma$, $\beta \circ \gamma$ with α, β being the same as in (1.3).

Corollary 4.3 *Let $\gamma : I \to M$ be a slant curve with null normal in a 3-dimensional α-Kenmotsu manifold M, for which $c^2 \neq 1$. Then $\alpha = \pm g(N, \varphi\dot\gamma) \in \mathbb{R}$ and*

$$\kappa = \alpha c, \quad N = -\alpha(\xi - c\dot\gamma \pm \varphi\dot\gamma), \quad W = \frac{-1}{2\alpha(1-c^2)}(\xi - c\dot\gamma \mp \varphi\dot\gamma).$$

In the particular case when γ is a Legendre curve, we have

$$\kappa = 0, \quad N = -\alpha(\xi \pm \varphi\dot\gamma), \quad W = \frac{-1}{2\alpha}(\xi \mp \varphi\dot\gamma).$$

where for simplicity we write α, β instead of the composed functions $\alpha \circ \gamma$, $\beta \circ \gamma$ with α, β being the same as in (1.3).

5 Examples illustrating theorems

Legendre curves with null normal that will appear in examples are from my previous paper [15].

Example 1 Let $\mathbb{R}^3$ be the Cartesian space and (x, y, z) be the Cartesian coordinates in it. Define the standard almost paracontact structure (φ, ξ, η) on $\mathbb{R}^3$ by

$$\varphi\partial_1 = \partial_2 - 2x\partial_3, \quad \varphi\partial_2 = \partial_1, \quad \varphi\partial_3 = 0, \quad \xi = \partial_3, \quad \eta = 2xdy + dz, \tag{5.1}$$

where $\partial_1 = \dfrac{\partial}{\partial x}$, $\partial_2 = \dfrac{\partial}{\partial y}$ and $\partial_3 = \dfrac{\partial}{\partial z}$. By certain direct calculations, one verifies that

$$[\varphi, \varphi](\partial_i, \partial_j) - 2d\eta(\partial_i, \partial_j)\xi = 0, \quad 1 \leqslant i < j \leqslant 3,$$

so that (1) is satisfied and the structure is normal.

Suppose that $M = \mathbb{R}^2 \times \mathbb{R}_+ \subset \mathbb{R}^3$ and consider a normal almost paracontact metric structure on M defined in the following way: (φ, ξ, η) is the structure (5.1) restricted to M and g is the Lorentz metric given by

$$[g(\partial_i, \partial_j)] = \begin{bmatrix} -2z & 0 & 0 \\ 0 & 4x^2 + 2z & 2x \\ 0 & 2x & 1 \end{bmatrix}.$$

For the Levi-Civita connection, we have

$$\nabla_{\partial_1}\partial_1 = -\frac{x}{z}\,\partial_2 + \Big(1 + \frac{2x^2}{z}\Big)\partial_3, \quad \nabla_{\partial_1}\partial_2 = \nabla_{\partial_2}\partial_1 = \frac{x}{z}\,\partial_2 + \Big(1 - \frac{2x^2}{z}\Big)\partial_3,$$
$$\nabla_{\partial_1}\partial_3 = \nabla_{\partial_3}\partial_1 = \nabla_{\partial_2}\partial_3 = \nabla_{\partial_3}\partial_2 = \frac{1}{2z}\,\partial_1 + \frac{1}{2z}\,\partial_2 - \frac{x}{z}\,\partial_3,$$
$$\nabla_{\partial_2}\partial_2 = \frac{2x}{z}\,\partial_1 + \frac{x}{z}\,\partial_2 - \Big(1 + \frac{2x^2}{z}\Big)\partial_3, \quad \nabla_{\partial_3}\partial_3 = 0.$$

Using the above and (1.2), we find $\alpha = \beta = (2z)^{-1}$.

$$(a) \qquad \gamma(t) = (0, -2\sqrt{-t}, t), \quad t < 0$$

is a Frenet slant curve. For such a curve $\varepsilon_1 = -1$, $c = 1$, $\alpha(\gamma(t)) = \beta(\gamma(t)) = \dfrac{1}{2t}$, $\delta(t) = \dfrac{1}{t}$, $\kappa = -\dfrac{\sqrt{5}}{\sqrt{2}t}$, $\tau = -\dfrac{3}{2t}$.

$$(b) \qquad \gamma(t) = (1/4, t, 3/8),$$

is a slant curve with null normal and $\varepsilon_1 = 1$.

For such a curve, we have $c = 1/2$, $\alpha(\gamma(t)) = 4/3$, $\beta(\gamma(t)) = 4/3$ and

$$T = (0, 1, 0), \quad N = (4/3, 2/3, -4/3)$$
$$W = (-1/2, 1/4, -1/2), \quad \kappa = -2/3.$$

$$(c) \quad \gamma(t) = (\sqrt{t}, -a\sqrt{t}, at), \quad t > 0, \quad a = \sqrt[3]{1-b} + \sqrt[3]{1+b}, \quad b = \sqrt{\frac{26}{27}},$$

is a Legendre curve with null normal and $\varepsilon_1 = 1$. For such a curve, we have $\alpha(\gamma(t)) = (2at)^{-1}$, $\beta(\gamma(t)) = (2at)^{-1}$, $\alpha(\gamma(t)) = g(N, \varphi\dot{\gamma}) = 1/(2at)$ and

$$T = ((2\sqrt{t})^{-1}, -a(2\sqrt{t})^{-1}, a), \quad N = (1/4t^{-3/2}, -1/(4a)t^{-3/2}, 0),$$
$$W = (-2a^2\sqrt{t}, 2a\sqrt{t}, -8at), \quad \kappa = \frac{1-2a}{2at}.$$

Example 2 Suppose that $M = \mathbb{R}^2 \times \mathbb{R}_+$. Define a normal almost paracontact structure (φ, ξ, η) on M by

$$\varphi\partial_1 = \partial_2, \quad \varphi\partial_2 = \partial_1, \quad \varphi\partial_3 = 0, \quad \xi = \partial_3, \quad \eta = dz, \tag{5.2}$$

and compatible with this structure a Lorentz metric

$$[g(\partial_i, \partial_j)] = \begin{bmatrix} -2z & 0 & 0 \\ 0 & 2z & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

The quadruple (φ, ξ, η, g) becomes a normal almost paracontact metric structure on M. For the Levi-Civita connection, we find

$$\begin{aligned} &\nabla_{\partial_1}\partial_1 = \partial_3, \quad \nabla_{\partial_1}\partial_2 = \nabla_{\partial_2}\partial_1 = 0, \quad \nabla_{\partial_1}\partial_3 = \nabla_{\partial_3}\partial_1 = \frac{1}{2z}\,\partial_1, \\ &\nabla_{\partial_2}\partial_2 = -\partial_3, \quad \nabla_{\partial_2}\partial_3 = \nabla_{\partial_3}\partial_2 = \frac{1}{2z}\,\partial_2, \quad \nabla_{\partial_3}\partial_3 = 0. \end{aligned}$$

Using the above and (1.2), we get $\alpha = (2z)^{-1}$ and $\beta = 0$.

The curve

$$\gamma(t) = \left(\cosh t, \sinh t, \frac{1}{2}\right),$$

is a Legendre curve with $\varepsilon_1 = 1$ and null normal in M.

Then for γ, we get $\alpha(\gamma(t)) = 1$, $\delta(t) = -1$ and

$$\begin{aligned} &T = (\sinh t, \cosh t, 0), \quad N = (\cosh t, \sinh t, -1), \\ &W = -\frac{1}{2}(\cosh t, \sinh t, 1), \quad \kappa = 0. \end{aligned}$$

References

[1] W. B. Bonnor, *Curves with null normals in Minkowski space-time.* In: A random walk in relativity and cosmology, Essays Hon. P. C. Vaidya and A. K. Raychaudhuri, pp. 33-47, Wiley Easten Limited (1985).

[2] B. Cappelletti Montano, *Bi-Legendrian structures and paracontact geometry*, Int. J. Geom. Meth. Mod. Phys. 6 (2009), 487-504.

[3] C. Călin, M. Crasmareanu, *Slant Curves in 3-dimensional Normal Almost Contact Geometry* Mediterr. J. Math., (2012)

[4] C. Călin, M. Crasmareanu, *Slant curves and particles in 3-dimensional warped products and their Lancret invariants*, Bull. Austr. Math. Soc., available on CJO2012. doi:10.1017/S0004972712000809.

[5] C. Călin, M. Crasmareanu, M. Munteanu, *Slant curves in 3-dimensional f -Kenmotsu manifolds* J. Math. Anal. Appl. **394** (2012), 400-407.

[6] P. Dacko, *On almost para–cosymplectic manifolds*, Tsukuba J. Math. **28** (2004), No. 1, 193-213.

[7] K. L. Duggal, *A Report on Canonical Null Curves and Screen Distributions for Lightlike Geometry* Acta Appl. Math. **95** (2007), 135149.

[8] K. L. Duggal and A. Bejancu, *Lightlike submanifolds of semi-Riemannian manifolds and applications.* Mathematics and its Applications, 364. Kluwer Academic Publishers Group, Dordrecht, 1996.

[9] K. L. Duggal and D. H. Jin, *Null curves and hypersurfaces of semi-Riemannian manifolds.* World Scientific Publishing, Singapore, 2007.

[10] S. Erdem, *On almost (para)contact (hyperbolic) metric manifolds and harmonicity of* (φ, φ')*-holomorphic maps between them*, Houston J. Math. **28** (2002), 21-45.

[11] K. İlarslan and E. Nešović, *Timelike and null normal curves in Minkowski space* $\mathbb{E}_1^3$, Indian J. Pure Appl. Math. **35** (2004), No. 7, 881-888.

[12] J.-I. Inoguchi, *Biharmonic curves in Minkowski 3-space. II*, Int. J. Math. Math. Sci. (2006), Art. ID 92349, 4 pp.

[13] H. Kocayigit and H. H. Hacisalihoglu, *1-type and biharmonic frenet curves in Lorentzian 3-space* Iranian Journal of Science and Technology, Transaction A, **33** (2009), No. A2

[14] S. Kaneyuki and F. L. Williams, *Almost paracontact and paraHodge structures on manifolds*, Nagoya Math. J. **99** (1985), 173-187.

[15] J. Wełyczko, *On Legendre curves in 3-dimensional normal almost paracontact metric manifolds*, Results Math. **54** (2009), 377-387.

[16] S. Zamkovoy, *Canonical connections on paracontact manifolds*, Ann. Glob. Anal. Geom. **36** (2009), No. 1, 37-60.